Sheet 1 of 9




FIGURE 1

Sheet 2 of 9





FIGURE 2

Sheet 3 of 9



Concentration of intracellular 30MG measured using radiolabeled 30MG


FIGURE 3

EST AVAILABLE COPY

Title: PRESERVATION OF BIOMATERIALS WITH TRANSPORTED PRESERVATION AGENTS By: Mehmet Toner et al.


Atty Docket No. 22727-138

Sheet 4 of 9



Percentage of dead cells after loading glucoses or DMSO


FIGURE 4



Metabolic activity was assessed measuring the MTT reduction activity


FIGURE 5

Sheet 5 of 9




Viability of cryopreserved mammalian cells with glucose loading

FIGURE 6



Viability of cryopreserved mammalian cells preserved using different agents


Sheet 6 of 9




Cell survival as a function of residual water in the sample after drying with or without 3OMG loading

By: Mehmet Toner et al. Atty Docket No. 22727-138


Sheet 7 of 9






Kinetics of 30MG uptake and efflux on hepatocytes and effects on viability and metabolic activity


Sheet 8 of 9




Post-thaw viability of cryopreserved hepatocytes

Sheet 9 of 9







Albumin production, urea production, and cytochrome P450 activity of treated hepatocytes

FIGURE 11